



组蛋白去乙酰化酶抑制剂调控缺血性脑卒中的分子机制*

潘冯缘^{1,2)} 徐怡薇^{1,2,3)} 邹歆好^{1,2)} 陈 静^{1,2)**}

(¹⁾ 三峡大学健康医学院, 宜昌 443002; (²⁾ 三峡大学国家中医药管理局中药药理科研三级实验室, 宜昌 443002;

(³⁾ 三峡大学现代药物制剂宜昌市重点实验室, 宜昌 443002)

摘要 缺血性脑卒中 (ischemic stroke, IS) 约占所有卒中病例的 80%, 是全球范围内导致死亡和长期残疾的主要神经系统疾病, 其核心病理机制为脑血流中断导致的神经细胞坏死和脑组织缺血性损伤, 涉及细胞凋亡、炎症反应及氧化应激等多重分子过程。组蛋白去乙酰化酶 (histone deacetylase, HDAC) 通过去除组蛋白上的乙酰基团 (acetyl group, Ac) 调控染色质构象与基因表达, 在神经细胞存活、炎症反应调控及血脑屏障稳态调节中发挥重要的作用。组蛋白去乙酰化酶抑制剂 (HDAC inhibitor, HDACi) 可抑制 HDAC 的活性, HDACi 不仅作用于组蛋白, 还可作用于热休克转录因子 1 (HSF1)、分子伴侣热休克蛋白 90 (HSP90) 及细胞骨架蛋白 α 微管蛋白 (α -tubulin) 在内的多种非组蛋白, 进而调控下游基因转录及相关信号通路, 在 IS 模型中表现出显著神经保护作用。本文系统综述 HDACs 的分类、HDACi 的作用机制, 及其在 IS 后细胞凋亡抑制、神经炎症调控、血脑屏障修复及认知功能改善中的多重效应, 同时探讨 HDACi 的靶向策略与临床应用前景, 为 HDACi 治疗 IS 提供了新的理论依据与科学的研究方向。

关键词 组蛋白去乙酰化酶, 组蛋白去乙酰化酶抑制剂, 缺血性脑卒中, 神经保护

中图分类号 R3, Q7

DOI: 10.3724/j.pibb.2025.0403

CSTR: 32369.14.pibb.20250403

缺血性脑卒中 (ischemic stroke, IS), 又称脑梗死, 是最常见的卒中类型, 占有卒中病例的 80% 以上^[1], 其核心病理机制始于脑血流中断, 引发脑组织缺氧及能量代谢障碍, 进而诱发神经元兴奋性毒性与钙离子超载, 导致线粒体损伤, 最终造成细胞凋亡与坏死。缺氧状态下, 自由基大量积累引发氧化应激与炎症反应, 炎症激活后破坏血脑屏障 (blood-brain barrier, BBB), 进一步诱发脑水肿并扩大缺血区域。从分子层面而言, IS 的发生与发展, 主要是细胞凋亡、神经炎症与氧化应激等多重损伤途径共同作用的结果^[2]。

组蛋白去乙酰化酶 (histone deacetylase, HDAC) 可特异性去除组蛋白上的乙酰基团, 使染色质结构趋于紧密, 阻碍转录因子与 RNA 聚合酶的结合, 最终实现基因沉默^[3] (图 1)。HDACs 在多种疾病的发生发展中发挥关键作用, 尤其在癌症、神经退行性疾病及心血管疾病中, 其异常表达或活性改变会导致基因表达失调, 进而引发病理性

变化, 因此抑制 HDACs 活性已成为重要的治疗策略^[4]。在神经科学领域, 抑制 HDACs 的作用已被证实具有神经保护效应, HDAC 抑制剂 (HDAC inhibitor, HDACi) 可阻断 HDACs 的去乙酰化功能, 维持或升高组蛋白乙酰化水平, 使染色质保持松散构象, 从而促进转录因子结合和基因转录激活^[5]。已经有研究证实 HDACi 对脑损伤具有显著神经保护作用^[6], 近年来相关研究证据表明 HDACi 可通过抑制卒中后神经细胞凋亡、缓解神经炎症、修复血脑屏障损伤及改善卒中后认知障碍 (post-stroke cognitive impairment, PSCI), 发挥 IS 治疗作用, 本文将系统综述 HDACi 调控 IS 的分子机制以及相关治疗研究进展。

* 湖北省自然科学基金 (2024AFD99) 资助项目。

** 通讯联系人。

Tel: 0717-6392029, E-mail: chenjing@ctgu.edu.cn

收稿日期: 2025-09-09, 接受日期: 2025-11-27




Fig. 1 Histone deacetylase and histone deacetylase inhibitors
图1 HDAC和HDAC抑制剂

1 HDAC的分类与结构

1.1 HDAC的分类

迄今为止, HDACs 家族已鉴定出 18 个成员。根据进化起源差异, HDACs 可大致分为 I、II、III、IV 四大类^[7]; 而依据催化方式不同, 其可分为两大类^[8]: 一类是锌离子 (Zn²⁺) 依赖型经典 HDACs, 包括 I 类 (HDAC1~3、HDAC8)、IIa 类 (HDAC4、5、7、9)、IIb 类 (HDAC6、10) 和 IV 类 (HDAC11); 另一类是烟酰胺腺嘌呤二核苷酸 (NAD⁺) 依赖型 III 类去乙酰化酶, 通常称为沉默信息调节因子 (sirtuins, SIRT1)。该分类既明确了各类 HDACs 在催化机制上的显著差异, 也暗示了它们在细胞代谢过程和表观遗传调控中具有不同的生物学功能。

1.2 HDAC的结构

在神经科学领域, I 类和 II 类 HDACs 因在中枢神经系统发育和功能中扮演关键角色, 成为研究最为广泛的亚型^[9]。I 类 HDACs 定位于细胞核, 属于广泛表达的核酶^[10]; II 类 HDACs 可在细胞质和细胞核之间作为信号转运蛋白, 通过调控基因表达、细胞周期、细胞分化等多个生理过程, 影响神经系统与免疫系统的功能^[11]。IIa 类 HDACs 作为信号转导因子, 其 N 端调控结构域包含 2 或 3 个保守丝氨酸残基, 这些残基能进行可逆性磷酸化修饰; 磷酸化可促进 14-3-3 蛋白结合, 进而介导 HDACs 的核出口并解除对其靶基因的抑制^[12]。III 类 HDACs 是区别于 I 类和 II 类的独立家族, 主要包括 SIRT1s,

其家族成员 SIRT1~7 广泛参与调控 DNA 修复、细胞凋亡、抗氧化反应以及代谢稳态等过程, 该类 HDACs 的活性依赖 NAD⁺, 其中 SIRT1 作为最具代表性的成员, 在神经退行性疾病和心血管疾病领域已有广泛研究^[13-14]。IV 类 HDAC 与 II 类在序列上有一定相似性, 目前仅发现 HDAC11, 相关研究仍较为匮乏。

2 HDAC抑制剂 (HDACi) 分类与功能

2.1 HDACi的分类

按化学结构分类, HDACi 大致可分为异羟肟酸类、短链脂肪酸类、苯甲酰胺类及其他类型^[15]。异羟肟酸类为广谱抑制剂^[16], 分子质量<500 u, 易穿过 BBB, 其含有的异羟肟酸基团 (-CONHOH) 可强效螯合锌离子, 竞争性占据 HDAC 催化活性中心, 抑制 I 类 HDACs (1/2/3/8)、IIb 类 HDACs (6/10) 和 IV 类 HDACs (11) 的活性^[17]。曲古抑菌素 A (Trichostatin A, TSA) 属于异羟肟酸类 HDACi, 对 7 日龄大鼠缺氧缺血模型, 损伤后给予 TSA 治疗, 可减少大鼠神经元及少突胶质细胞祖细胞丢失, 同时促进脑源性神经营养因子 (brain-derived neurotrophic factor, BDNF) 表达并激活 TrkB 受体^[18]; Tubacin 能减少促炎细胞因子介导的内皮一氧化氮合酶 (endothelial nitric oxide synthase, eNOS) 降解, 改善小鼠心血管内皮功能, 其通过上调 eNOS 表达可能对内皮功能障碍相关的心血管疾病具有治疗潜力^[19]; 伏立诺他 (Vorinostat Zolinza, SAHA) 属于广谱抑制剂, 能

够降低白介素 (interleukin, IL) -1 β 、IL-6、肿瘤坏死因子 α (tumor necrosis factor- α , TNF- α) 等促炎因子的水平^[20]。短链脂肪酸类主要抑制 I 类 HDACs (1/2/3) 和 IIa 类 HDACs (4/5/7/9), 具有一定靶向选择性, 主要由肠道产生^[21], 代表药物为丙戊酸钠 (sodium valproate, VPA) 和丁酸钠 (sodium butyrate, SB)。研究发现, 在小鼠大脑中动脉闭塞 (MCAO) 模型中, SB 可下调 TNF- α 和一氧化氮合酶 (nitric oxide synthase 2, NOS2)、上调 IL-10, 通过小胶质细胞介导的表观遗传调节发挥神经保护作用并缓解炎症反应^[22]。苯甲酰胺

类 HDACi 的代表药物包括恩替诺特 (Entinostat) 和恩替司他 (MS-275), 常作为 I 类 HDACi, 靶向作用于 HDAC1~3 和 HDAC11^[23], 研究显示, MS-275 在缺血再灌注模型中可有效缩小脑梗死面积, 改善神经功能评估结果^[24]。其他类型中, β 萘酚衍生物类、噻吩并嘧啶酮衍生物类主要抑制 III 类去乙酰化酶^[25-26], HDAC11 特异性抑制剂 PB94 包含苯乙烯基 (styryl) 和苯甲酰胺 (Benzamide) 两大核心基团, 通过特定连接形成独特的三环骨架^[27] (表 1)。

Table 1 Classification, representative drugs and mechanisms of HDAC inhibitors

表 1 HDACi 的分类、代表药物以及作用机制

化学类别	代表药物	作用机制
异羟肟酸类	TSA	抑制 I 类、II 类、IV 类 HDAC ^[18]
	Tubacin	高度选择性抑制 HDAC6 ^[19]
	SAHA	抑制 I 类、II 类、IV 类 HDAC ^[20]
短链脂肪酸类	SB	抑制 I 类和 IIa 类 HDAC ^[21]
	VPA	
苯甲酰胺类	Entinostat	抑制 I 类 HDAC ^[23-24]
	MS-275	
	RGFP966	
β 萘酚衍生物	Sirtinol	主要抑制 HDAC1/2/3 ^[28]
噻吩并嘧啶酮衍生物	AGK2	选择性抑制 SIRT1 和 SIRT2 ^[25]
		选择性抑制 SIRT2 ^[26]
苯乙烯基苯甲酰胺	PB94	选择性抑制 IV 类 HDAC ^[27]

TSA: 曲古抑菌素 A (Trichostatin A); SAHA: 伏立诺他 (Vorinostat Zolinza); SB: 丁酸钠 (sodium butyrate); VPA: 丙戊酸钠 (sodium valproate); Entinostat: 恩替诺特; MS-275: 恩替司他; Sirtinol: 西尔替诺; AGK2: SIRT2 抑制剂; SIRT1: 沉默信息调节因子 1 (sirtuin1); SIRT2: 沉默信息调节因子 2 (sirtuin2)。

2.2 HDACi 的功能

HDACi 作为一类重要的表观遗传修饰剂, 其经典机制是通过抑制 HDACs 活性升高组蛋白乙酰化水平, 进而松弛染色质并激活 *p21* 等基因转录进程^[29], 此外, HDACi 的作用范畴不限于组蛋白, 还可广泛作用于转录因子热休克因子 1 (heat shock factor 1, HSF1)、分子伴侣 HSP90 及细胞骨架蛋白 α 微管蛋白 (α -tubulin) 在内的多种非组蛋白底物^[30], 通过调控这些底物的乙酰化状态影响细胞功能, 同时激活抑癌基因、神经营养因子等相关基因转录^[31]。不仅如此, HDACi 还能调控 p53、核因子 κ B (nuclear factor κ B, NF- κ B) 等 50 余种非组蛋白底物的乙酰化修饰, 对细胞凋亡、炎症反应、代谢调控及细胞骨架稳定性等关键通路进行重编程^[32-33]。

在神经疾病领域, HDACi 可通过抑制神经炎

症、抵抗氧化应激、抑制细胞凋亡和促进突触可塑性等途径发挥神经保护作用。HDAC2 抑制剂 CAY10683 能够有效抑制脂多糖 (lipopolysaccharide, LPS) 激活的 BV2 小胶质细胞, 以及 LPS 诱导的小鼠神经炎症模型中炎症细胞因子 TNF- α 和 IL-1 β 的表达^[34]; 研究发现^[35], IS 后, HDAC2 与凋亡神经元存在共定位关系; 另有研究证实^[36], HDACi 可通过抑制 NF- κ B 通路减少促炎因子分泌, 从而减轻 BBB 的炎症损伤。

IS 的传统治疗手段 (如溶栓、取栓等) 虽能在短时间内恢复脑部供血, 但易引发脑缺血再灌注损伤, 且存在出血风险高、救治时效性要求严苛等弊端^[37]。相比之下, HDACi 具备多靶点神经保护特性, 不仅能保护神经元, 还可维护 BBB 完整性, 降低出血风险^[6]。

3 HDACi调控IS的效应

3.1 HDACi调控IS后的细胞凋亡过程

IS会诱导细胞凋亡, 而激活内源性抗凋亡通路是保护神经元免受损伤的关键机制。HDACi可通过多途径干预这一过程, 进而激活抗凋亡通路发挥神经保护作用^[5]。有研究表明, HDACi能显著改善脑缺血后神经功能损伤, 减小脑梗死体积, 并且延缓神经细胞凋亡进程^[38]。

HDACi能促进组蛋白乙酰化, 使该基因启动子区域开放 (如p21/CIP1), 引发细胞周期阻滞, 从而阻止受损神经元启动凋亡程序^[39]; HDACi如VPA和SAHA通过增加组蛋白乙酰化水平, 下调促凋亡蛋白, 如caspase-3等, 同时上调抗凋亡蛋白, 如Bcl-2等, 从而抑制神经元凋亡^[40]。此外, 丁酸盐通过乙酰化HSF1促进热休克蛋白70 (heat shock protein 70, Hsp70) 表达, 维持细胞内蛋白质稳态, 抵抗内质网应激诱发的细胞凋亡^[41]。在促凋亡信号抑制方面, HDACi同样发挥重要作用。在p53通路中, 抑制HDAC1可诱导p53乙酰化, 阻断其与DNA的结合能力, 减少促凋亡基因的转录; 在JNK通路中, 抑制HDAC3能阻断Jun激酶 (Jun kinase, JNK) 磷酸化级联反应, 降低减少c-Jun氨基端激酶 (c-Jun N-terminal kinase, c-Jun) 的激活水平, 抑制应激诱导的神经元凋亡^[42]; HDAC6抑制剂 (如Tubastatin A) 可恢复 α -tubulin功能, 促进自噬清除受损线粒体, 同时阻断caspase-3激活, 进一步抑制凋亡进程^[43]。通过上述机制的协同作用, HDACi能有效调节细胞周期, 抑制神经元凋亡, 发挥神经保护作用。

3.2 HDACi调控IS后的神经炎症

IS会诱发神经炎症反应, 而HDACi通过表观遗传调控与信号通路干预双重作用, 在抑制该炎症进程中发挥关键作用, 核心机制包括沉默促炎基因、促进抗炎表型转换及调控炎症相关通路。在沉默促炎基因与抑制炎症体激活方面, HDAC3抑制剂RGFP966可激活Nrf2通路, 促进其核转位并激活下游交替氧化酶 (alternative oxidase, AO) 表达, 进而显著抑制高迁移率族蛋白B1 (high mobility group box 1, HMGB1) /Toll样受体4 (Toll-like receptor 4, TLR4) 介导的NOD样受体家族Pyrin域包含蛋白3 (NOD-like receptor protein 3) 炎症体激活^[44]; 在缺血再灌注损伤中, SAHA通过抑制NF- κ B通路减少促炎因子 (如IL-1 β 、IL-6、

TNF- α) 释放, 并激活核因子E2相关因子2 (nuclear factor erythroid-2-related factor 2) /血红素加氧酶1信号轴 (heme oxygenase-1) 抗氧化通路^[45]; IS还会诱导小胶质细胞中HDAC3过表达, 后者通过去乙酰化作用促进P65核易位, 诱导环磷酸鸟苷-磷酸腺苷合成酶 (cyclic GMP-AMP synthase, cGAS) 表达, 最终激活干扰素基因刺激蛋白 (stimulator of interferon genes, STING) /干扰素调节因子 (interferon regulatory factor, IRF) 通路, 反之, 敲除HDAC3可阻断该通路激活, 从而下调缺血诱导的脑组织损伤和神经炎症^[46]。此外, IV类HDACs (HDAC11) 通过抑制抗炎因子IL-10维持促炎微环境, 其选择性抑制剂PB94已被证实能有效阻断其促炎作用, 不过其具体抗炎机制仍有待进一步明确^[47]。在促进抗炎表型转换与调控细胞极化方面, HDAC3在巨噬细胞中可通过非去乙酰化依赖性机制激活TNF- α 、IL-6等促炎基因表达, 基于蛋白质靶向降解嵌合体 (proteolysis targeting chimera, PROTAC) 技术开发的HDAC3降解剂P7, 能有效阻断M0型巨噬细胞向促炎表型M1极化, 显著减少促炎因子分泌并增加IL-10等抗炎因子释放^[48]。

HDAC2通过调控促炎基因的染色质重塑参与炎症调控, 其失活会导致促炎基因表达升高^[49]; IIa类HDACs (HDAC4/5) 主要促进脑卒中后星形胶质细胞活化及瘢痕形成^[50], 而抑制HDAC4/5可降低STAT3磷酸化水平, 抑制星形胶质细胞的反应性胶质增生^[51], 因此IIa类HDACi更适合在卒中修复期使用。综上, HDACi通过表观遗传调控和信号通路干预双重作用, 发挥对神经炎症的抑制作用, 尤其是针对HDAC3的靶向抑制或降解, 揭示了精准调控神经炎症的新方向, 为IS的治疗提供了新思路。

3.3 HDACi调控IS后的BBB损伤

IS后, BBB会发生多阶段损伤, 其破坏过程分为急性期、亚急性期和慢性期^[52]。急性期时, 缺氧导致ATP耗竭, 使Na⁺/K⁺-ATP酶功能失效, 引发细胞毒性水肿, 同时闭合蛋白 (occludin)、密封蛋白5 (claudin-5) 等紧密连接蛋白被降解, 微血管通透性增加, 进而导致血管源性水肿。亚急性期伴随炎症反应级联激活, 小胶质细胞释放TNF- α 、IL-1 β 等炎症因子损伤内皮细胞, 加之白细胞浸润和活性氧类 (reactive oxygen species, ROS) 大量释放, 进一步破坏BBB完整性, 增加出血转化风

险^[53]。慢性期则以周细胞凋亡、基底膜降解为主要特征,导致微血管稳定性丧失、血管再生受阻,最终延迟脑组织修复进程^[54]。

HDACi通过多种机制保护或修复BBB,包括抗炎抗氧化、调控紧密连接蛋白表达、抑制异常通路激活及促进修复相关基因表达。HDAC3抑制剂可通过抑制炎症和氧化应激减轻BBB损伤,这一作用在缺血性脑损伤、阿尔茨海默病等疾病模型中已得到验证^[55]。在脑梗死缺血再灌注模型中,VPA通过抑制NF- κ B核转位和基质金属蛋白酶9(matrix metalloproteinase 9, MMP-9)表达,能够保护BBB完整性^[56]。有研究表明,益母草碱(leonurine, SCM-198)在氧糖剥夺/再灌注(OGD/R)模型中,通过抑制HDAC2加速occludin和claudin-5的恢复,降低内皮细胞单层通透性^[57];W2A-16通过激活Wnt/ β -catenin信号通路和抑制I类HDACs,增强跨内皮电阻(transendothelial electrical resistance, TEER),并上调BBB相关蛋白(如P-糖蛋白)的表达^[58]。

需要注意的是,HDACi作用于BBB的具体效应取决于疾病类型和使用时机。例如,在脑肿瘤治疗中,HDACi可能通过暂时性开放BBB促进药物输送^[59],而在脑缺血治疗中则以保护BBB完整性为主要作用。

3.4 HDACi调控IS后的认知障碍

IS后,PSCI是临床常见后遗症^[60],但目前针对性治疗手段仍较为有限。HDACs在PSCI的病理进程中发挥关键调控作用,不同亚型的功能具有显著特异性,为HDACi的干预提供了明确靶点。I类HDACs主要定位于细胞核^[61],研究证实其在卒中后表达显著上调,是加剧神经元损伤、抑制神经可塑性及损害认知功能的核心因素^[62],因此成为HDACi治疗PSCI的主要靶点。IIa类HDACs可在核质间穿梭,其中HDAC4常被用作PSCI的潜在标志物^[63],主要参与记忆功能与认知障碍的调控。研究发现^[64],IS后PSCI患者的HDAC4表达水平低于仅发生卒中而无认知障碍的患者,这一特征可作为PSCI早期筛查的参考标准。动物实验显示,IS 4周后,大鼠海马区HDACs活性增强,导致认知功能受损,而应用I类HDACi SB可有效降低海马区HDACs活性,提高组蛋白乙酰化水平,促进神经可塑性相关基因表达,进而改善认知功能障碍^[65]。

HDACi通过多种机制协同改善PSCI的病理过

程。其一,促进组蛋白乙酰化,解除神经营养因子(如BDNF)和突触相关基因的转录抑制,激活TrkB受体及其下游信号通路,改善神经元存活与突触功能,提升学习记忆能力^[66-67];其二,抑制NF- κ B、转录激活因子1(activator protein-1, AP-1)等炎症通路,减少促炎因子表达^[68],改善炎症微环境,发挥神经保护作用^[69];其三,增强抗氧化防御能力,清除ROS,减轻氧化应激对神经元的损伤^[70],通过调节线粒体相关蛋白的乙酰化状态,促进线粒体生物发生,提升电子传递链效率,减轻氧化应激并优化脑内能量供应^[71];其四,通过激活磷脂酰肌醇3激酶(phosphatidylinositol 3 kinase)/蛋白激酶B(protein kinase B)/糖原合成酶激酶3 β (glycogen synthase kinase 3 beta)通路发挥抗凋亡作用,保护海马和皮层等认知关键脑区的神经元^[72];其五,促进海马齿状回神经发生,修复损伤的神经元网络,改善空间学习和记忆能力^[73];其六,针对HDAC6的特异性抑制可增加 α -tubulin乙酰化水平,改善轴突运输效率,减少tau蛋白异常聚集,同时增强热休克蛋白功能,保护神经元结构和功能完整性^[74]。综上,HDACs在IS后认知障碍的发生发展中发挥关键调控作用。当前,HDACi治疗PSCI的相关研究持续深入,其多机制协同的干预特性,为PSCI的临床治疗开辟了新路径,展现出广阔的应用前景。

4 总结与展望

HDACi可通过调节组蛋白乙酰化水平,改善神经细胞功能及生存微环境,减轻缺血性脑卒中后的脑损伤,推动神经功能恢复^[75]。作为极具潜力的神经保护治疗手段,HDACi已在多项研究中证实对神经系统疾病的干预价值,其机制主要涉及乙酰化修饰调控。一方面通过增强组蛋白乙酰化,激活神经营养因子、Hsp70、Bcl-2等促存活、修复相关基因表达,同时抑制NLRP3等促炎相关基因活性;另一方面通过上调细胞内乙酰化水平,调控Nrf2、NF- κ B、BDNF/TrkB等信号通路,实现抑制细胞凋亡、减少炎症反应、修复BBB损伤、缓解PSCI等协同效应,最终促进神经细胞存活与损伤修复(图2)。目前,临床常用的HDACi包括VPA、SB、TSA、Entinostat、SAHA等,其中VPA是最具代表性的药物之一,已应用于多种神经系统疾病的临床治疗。临床数据显示,VPA具有脂溶性,能够穿透BBB进入脑组织,在卒中治疗中,




Fig. 2 The mechanism of HDACi in treating ischemic stroke
图2 HDACi治疗缺血性脑卒中的机制

HDACi: 组蛋白去乙酰化酶抑制剂 (histone deacetylases inhibitors); p21: 细胞周期依赖性蛋白激酶抑制因子1A (cyclin-dependent kinase inhibitor 1A); CIP1: 周期素依赖激酶相互作用蛋白1 (cyclin-dependent kinase interacting protein 1); Bcl-2: B细胞淋巴瘤/白血病-2基因 (B-cell lymphoma-2); HSF1: 热休克转录因子1 (heat shock transcription factor 1); p53: 磷酸化蛋白53 (phosphorylated protein 53); JNK: Jun激酶 (Jun kinase); c-Jun: c-Jun氨基端激酶 (c-Jun N-terminal kinase); BDNF: 脑源性神经营养因子 (brain-derived neurotrophic factor); TrkB: 神经营养酪氨酸激酶受体2型 (neurotrophic tyrosine kinase receptor, type 2); PIK: 磷酸肌醇3激酶 (phosphoinositide-3-kinase); Akt: 蛋白激酶B (protein kinase B); GSK3β: 糖原合成酶激酶3β (recombinant glycogen synthase kinase 3 beta); α-tubulin: α微管蛋白; tau: 微管相关蛋白 (microtubule-associated protein tau); NF-κB: 核因子κB (nuclear factor κB); AP-1: 活化蛋白1 (activator protein-1); ROS: 活性氧类 (reactive oxygen species); occludin: 闭合蛋白; claudin-5: 密封蛋白5; MMP-9: 基质金属蛋白酶9 (matrix metalloproteinase 9); β-catenin: β连环蛋白; TEER: 跨内皮电阻 (transendothelial electrical resistance); Nrf2: 核因子E2相关因子2 (nuclear factor erythroid-derived 2-related factor 2); AO: 下游交替氧化酶 (alternative oxidase); NLRP3: 核苷酸结合寡聚化结构域样受体蛋白3 (NOD-like receptor protein 3); STAT3: 信号转导与转录激活因子3 (signal transducer and activator of transcription 3); HO-1: 血红素氧合酶1 (heme oxygenase-1); TNF-α: 肿瘤坏死因子α (tumor necrosis factor-α); IL-1β: 白介素-1β (interleukin-1β)。

剂量 100~250 mg/kg 的短期应用 (如 7~28 d) 的耐受性良好 [76]。

尽管 HDACi 在神经损伤及神经退行性疾病治疗中表现出显著潜力, 但临床转化过程中仍面临关键挑战, 长时间使用 HDACi 易引发毒副作用, 且可能影响正常细胞功能, 这在一定程度上限制了其广泛应用。以临床已应用的 VPA 为例, 其常见副作用包括胃肠道反应、头晕、恶心等, 严重时还可导致肝毒性, 需重点关注用药安全性 [76]。因此, 研发高选择性、低毒性的新型 HDACi 药物成为未

来研究的关键要点。此外, VPA 的临床潜力需进一步通过随机对照试验验证, 尤其是针对复发性卒中预防和神经修复的优化方案, 如联合细胞疗法或靶向 HDACi [77]。

HDACi 在神经系统疾病治疗领域具有广阔的应用潜力, 尤其在 IS 的防治中已表现出良好的临床转化前景。未来研究需进一步探讨 HDACi 的作用机制, 同时基于精准靶向理论与技术筛选最优治疗方案。针对 HDACi 临床应用中面临的不良反应控制、给药剂量优化等核心问题, 进一步提升其靶

向选择性、优化药物递送系统是关键研究方向。随着表观遗传调控技术、药物研发技术及递送系统的创新突破, HDACi有望发展为治疗神经损伤及神经退行性疾病的高效、安全干预手段, 为相关疾病的临床诊疗提供新的思路。

参 考 文 献

- [1] Sun L R, Lynch J K. Advances in the diagnosis and treatment of pediatric arterial ischemic stroke. *Neurotherapeutics*, 2023, **20**(3): 633-654
- [2] Ho J P, Powers W J. Contemporary management of acute ischemic stroke. *Annu Rev Med*, 2025, **76**(1): 417-429
- [3] Li J, Yan X, Liang C, *et al.* Comprehensive analysis of the differential expression and prognostic value of histone deacetylases in glioma. *Front Cell Dev Biol*, 2022, **10**: 840759
- [4] OuYang C, Shu G, Liu J, *et al.* HDAC5, negatively regulated by miR-148a-3p, promotes colon cancer cell migration. *Cancer Sci*, 2022, **113**(8): 2560-2574
- [5] Sixto-López Y, Correa-Basurto J. HDAC inhibition as neuroprotection in COVID-19 infection. *Curr Top Med Chem*, 2022, **22**(16): 1369-1378
- [6] Hummel R, Dorochow E, Zander S, *et al.* Valproic acid treatment after traumatic brain injury in mice alleviates neuronal death and inflammation in association with increased plasma lysophosphatidylcholines. *Cells*, 2024, **13**(9): 734
- [7] Wu H, Yin X, Zhao X, *et al.* HDAC11 negatively regulates antifungal immunity by inhibiting Nos2 expression *via* binding with transcriptional repressor STAT3. *Redox Biol*, 2022, **56**: 102461
- [8] Boateng A T, Abaidoo-Myles A, Bonney E Y, *et al.* Isoform-selective versus nonselective histone deacetylase inhibitors in HIV latency reversal. *AIDS Res Hum Retroviruses*, 2022, **38**(8): 615-621
- [9] Turkman N, Xu S, Huang C H, *et al.* High-contrast PET imaging with [(18)F] NT160, a class-IIa histone deacetylase probe for *in vivo* imaging of epigenetic machinery in the central nervous system. *J Med Chem*, 2023, **66**(8): 5611-5621
- [10] Kuang J, Li P, Zhai Z, *et al.* Exclusion of HDAC1/2 complexes by oncogenic nuclear condensates. *Mol Cancer*, 2024, **23**(1): 85
- [11] Broide R S, Redwine J M, Aftahi N, *et al.* Distribution of histone deacetylases 1-11 in the rat brain. *J Mol Neurosci*, 2007, **31**(1): 47-58
- [12] Martin M, Kettmann R, Dequiedt F. Class IIa histone deacetylases: conducting development and differentiation. *Int J Dev Biol*, 2009, **53**(2/3): 291-301
- [13] Zhang L, Li X, Chen Y, *et al.* Discovery of N-(2-aminophenyl)-4-(bis(2-chloroethyl)amino)benzamide as a potent histone deacetylase inhibitor. *Front Pharmacol*, 2019, **10**: 957
- [14] Grozinger C M, Chao E D, Blackwell H E, *et al.* Identification of a class of small molecule inhibitors of the sirtuin family of NAD-dependent deacetylases by phenotypic screening. *J Biol Chem*, 2001, **276**(42): 38837-38843
- [15] Oltra S S, Cejalvo J M, Tormo E, *et al.* HDAC5 inhibitors as a potential treatment in breast cancer affecting very young women. *Cancers (Basel)*, 2020, **12**(2): 412
- [16] Faraco G, Pancani T, Formentini L, *et al.* Pharmacological inhibition of histone deacetylases by suberoylanilide hydroxamic acid specifically alters gene expression and reduces ischemic injury in the mouse brain. *Mol Pharmacol*, 2006, **70**(6): 1876-1884
- [17] Bali P, George P, Cohen P, *et al.* Superior activity of the combination of histone deacetylase inhibitor LAQ824 and the FLT-3 kinase inhibitor PKC412 against human acute myelogenous leukemia cells with mutant FLT-3. *Clin Cancer Res*, 2004, **10**(15): 4991-4997
- [18] Zalewska T, Jaworska J, Sypecka J, *et al.* Impact of a histone deacetylase inhibitor-trichostatin A on neurogenesis after hypoxia-ischemia in immature rats. *Int J Mol Sci*, 2020, **21**(11): 3808
- [19] Chen J, Zhang J, Shaik N F, *et al.* The histone deacetylase inhibitor tubacin mitigates endothelial dysfunction by up-regulating the expression of endothelial nitric oxide synthase. *J Biol Chem*, 2019, **294**(51): 19565-19576
- [20] Çelik Z B, Günaydın C. Effects of suberoylanilide hydroxamic acid (SAHA) on the inflammatory response in lipopolysaccharide-induced N9 microglial cells. *Cureus*, 2022, **14**(12): e32428
- [21] Di Fede E, Ottaviano E, Grazioli P, *et al.* Insights into the role of the microbiota and of short-chain fatty acids in Rubinstein-taybi syndrome. *Int J Mol Sci*, 2021, **22**(7): 3621
- [22] Patnala R, Arumugam T V, Gupta N, *et al.* HDAC inhibitor sodium butyrate-mediated epigenetic regulation enhances neuroprotective function of microglia during ischemic stroke. *Mol Neurobiol*, 2017, **54**(8): 6391-6411
- [23] Batlevi C L, Kasamon Y, Bociek R G, *et al.* ENGAGE-501: phase II study of entinostat (SNDX-275) in relapsed and refractory Hodgkin lymphoma. *Haematologica*, 2016, **101**(8): 968-975
- [24] Murphy S P, Lee R J, McClean M E, *et al.* MS-275, a class I histone deacetylase inhibitor, protects the p53-deficient mouse against ischemic injury. *J Neurochem*, 2014, **129**(3): 509-515
- [25] Nikseresht S, Khodagholi F, Ahmadiani A. Protective effects of ex-527 on cerebral ischemia-reperfusion injury through necroptosis signaling pathway attenuation. *J Cell Physiol*, 2019, **234**(2): 1816-1826
- [26] Hernández-Jiménez M, Hurtado O, Cuartero M I, *et al.* Silent information regulator 1 protects the brain against cerebral ischemic damage. *Stroke*, 2013, **44**(8): 2333-2337
- [27] Chan A M, Mitchell A, Grogan L, *et al.* Histone deacetylase (HDAC) inhibitor specificity determinants are preserved in a class of dual HDAC/non-covalent proteasome inhibitors. *Bioorg Med Chem*, 2024, **104**: 117680
- [28] Chen S, Ye J, Wu G, *et al.* Histone deacetylase 3 inhibition ameliorates microglia-mediated neuro-inflammation *via* the SIRT1/Nrf2 pathway after traumatic spinal cord injury. *Neurorehabil Neural Repair*, 2023, **37**(8): 503-518
- [29] Whitmore M A, Li H, Lyu W, *et al.* Epigenetic regulation of host

- defense peptide synthesis: synergy between histone deacetylase inhibitors and DNA/histone methyltransferase inhibitors. *Front Immunol*, 2022, **13**: 874706
- [30] Sprinzen L, Garcia F, Mela A, *et al*. EZH2 inhibition sensitizes IDH1R132H-mutant gliomas to histone deacetylase inhibitor. *Cells*, 2024, **13**(3): 219
- [31] Soboleva S, Kurita R, Ek F, *et al*. Identification of potential chemical compounds enhancing generation of enucleated cells from immortalized human erythroid cell lines. *Commun Biol*, 2021, **4**(1): 677
- [32] Hu C, Peng K, Wu Q, *et al*. HDAC1 and 2 regulate endothelial VCAM-1 expression and atherogenesis by suppressing methylation of the GATA6 promoter. *Theranostics*, 2021, **11**(11): 5605-5619
- [33] Spange S, Wagner T, Heinzel T, *et al*. Acetylation of non-histone proteins modulates cellular signalling at multiple levels. *Int J Biochem Cell Biol*, 2009, **41**(1): 185-198
- [34] Jiao F Z, Wang Y, Zhang H Y, *et al*. Histone deacetylase 2 inhibitor CAY10683 alleviates lipopolysaccharide induced neuroinflammation through attenuating TLR4/NF- κ B signaling pathway. *Neurochem Res*, 2018, **43**(6): 1161-1170
- [35] Demyanenko S V, Dzreyan V A, Neginskaya M A, *et al*. Expression of histone deacetylases HDAC1 and HDAC2 and their role in apoptosis in the penumbra induced by photothrombotic stroke. *Mol Neurobiol*, 2020, **57**(1): 226-238
- [36] Lu Y, Chen Y, Xu S, *et al*. HDAC inhibitor attenuates rat traumatic brain injury induced neurological impairments. *Heliyon*, 2023, **9**(8): e18485
- [37] Liu J, Guo X, Yang L, *et al*. Effect of celastrol on lncRNAs and mRNAs profiles of cerebral ischemia-reperfusion injury in transient middle cerebral artery occlusion mice model. *Front Neurosci*, 2022, **16**: 889292
- [38] Guo M, Ye Y, Li X, *et al*. Kirenol alleviates cerebral ischemic injury by promoting synaptic plasticity via HDAC2-mediated BDNF expression. *Phytomedicine*, 2025, **143**: 156605
- [39] Buglio D, Georgakis G V, Hanabuchi S, *et al*. Vorinostat inhibits STAT6-mediated TH2 cytokine and TARC production and induces cell death in Hodgkin lymphoma cell lines. *Blood*, 2008, **112**(4): 1424-1433
- [40] Hunsberger J G, Fessler E B, Wang Z, *et al*. Post-insult valproic acid-regulated microRNAs: potential targets for cerebral ischemia. *Am J Transl Res*, 2012, **4**(3): 316-332
- [41] Sun J, Wu Q, Sun H, *et al*. Inhibition of histone deacetylase by butyrate protects rat liver from ischemic reperfusion injury. *Int J Mol Sci*, 2014, **15**(11): 21069-21079
- [42] Lo Cascio C, McNamara J B, Melendez E L, *et al*. Nonredundant, isoform-specific roles of HDAC1 in glioma stem cells. *JCI Insight*, 2021, **6**(17): e149232
- [43] Kaliszczak M, Trousil S, Ali T, *et al*. AKT activation controls cell survival in response to HDAC6 inhibition. *Cell Death Dis*, 2016, **7**(6): e2286
- [44] Xu L, An T, Jia B, *et al*. Histone deacetylase 3-specific inhibitor RGFP966 attenuates oxidative stress and inflammation after traumatic brain injury by activating the Nrf2 pathway. *Burns Trauma*, 2024, **12**: tkad062
- [45] Ling L, Zhou G, Zhang X, *et al*. A novel histone deacetylase inhibitor protects the blood-brain barrier by regulating NF- κ B and Nrf2 signaling pathways in OGD/R injury. *Arch Gerontol Geriatr*, 2025, **131**: 105739
- [46] Liao Y, Cheng J, Kong X, *et al*. HDAC3 inhibition ameliorates ischemia/reperfusion-induced brain injury by regulating the microglial cGAS-STING pathway. *Theranostics*, 2020, **10**(21): 9644-9662
- [47] Bai P, Liu Y, Yang L, *et al*. Development and pharmacochemical characterization discover a novel brain-permeable HDAC11-selective inhibitor with therapeutic potential by regulating neuroinflammation in mice. *J Med Chem*, 2023, **66**(23): 16075-16090
- [48] Zhao C, Chen S, Chen D, *et al*. Histone deacetylase 3-directed PROTACs have anti-inflammatory potential by blocking polarization of M0-like into M1-like macrophages. *Angew Chem Int Ed*, 2023, **62**(42): e202310059
- [49] van der Lugt T, Weseler A R, Vrolijk M F, *et al*. Dietary advanced glycation endproducts decrease glucocorticoid sensitivity *in vitro*. *Nutrients*, 2020, **12**(2): 441
- [50] Main P, Tan W J, Wheeler D, *et al*. Increased abundance of nuclear HDAC4 impairs neuronal development and long-term memory. *Front Mol Neurosci*, 2021, **14**: 616642
- [51] Yang H, Lan P, Hou Z, *et al*. Histone deacetylase inhibitor SAHA epigenetically regulates miR-17-92 cluster and MCM7 to upregulate MICA expression in hepatoma. *Br J Cancer*, 2015, **112**(1): 112-121
- [52] Gong S, Ma H, Zheng F, *et al*. Inhibiting YAP in endothelial cells from entering the nucleus attenuates blood-brain barrier damage during ischemia-reperfusion injury. *Front Pharmacol*, 2021, **12**: 777680
- [53] Zeng P, Wang X M, Ye C Y, *et al*. Mechanistic insights into the antidepressant effect of emodin: an integrated systems pharmacology study and experimental validation. *Aging*, 2021, **13**(11): 15078-15099
- [54] Huang J, Zhang R, Wang S, *et al*. Methamphetamine and HIV-tat protein synergistically induce oxidative stress and blood-brain barrier damage via transient receptor potential melastatin 2 channel. *Front Pharmacol*, 2021, **12**: 619436
- [55] Lu H, Ashiqueali R, Lin C I, *et al*. Histone deacetylase 3 inhibition decreases cerebral edema and protects the blood-brain barrier after stroke. *Mol Neurobiol*, 2023, **60**(1): 235-246
- [56] Wang Z, Leng Y, Tsai L K, *et al*. Valproic acid attenuates blood-brain barrier disruption in a rat model of transient focal cerebral ischemia: the roles of HDAC and MMP-9 inhibition. *J Cereb Blood Flow Metab*, 2011, **31**(1): 52-57
- [57] Zhang Q Y, Wang Z J, Sun D M, *et al*. Novel therapeutic effects of leonurine on ischemic stroke: new mechanisms of BBB integrity. *Oxid Med Cell Longev*, 2017, **2017**: 7150376

- [58] Inoue Y, Ren Y, Zhang S, *et al.* A novel histone deacetylase inhibitor W2A-16 improves the barrier integrity in brain vascular endothelial cells. *Front Cell Neurosci*, 2024, **18**: 1368018
- [59] Vazana U, Veksler R, Pell G S, *et al.* Glutamate-mediated blood-brain barrier opening: implications for neuroprotection and drug delivery. *J Neurosci*, 2016, **36**(29): 7727-7739
- [60] Zhang M S, Liang J H, Yang M J, *et al.* Low serum superoxide dismutase is associated with a high risk of cognitive impairment after mild acute ischemic stroke. *Front Aging Neurosci*, 2022, **14**: 834114
- [61] Cho E, Rowan-Carroll A, Williams A, *et al.* Development and validation of the TGx-HDACi transcriptomic biomarker to detect histone deacetylase inhibitors in human TK6 cells. *Arch Toxicol*, 2021, **95**(5): 1631-1645
- [62] Lin Y H, Dong J, Tang Y, *et al.* Opening a new time window for treatment of stroke by targeting HDAC2. *J Neurosci*, 2017, **37**(28): 6712-6728
- [63] Linares A, Assou S, Lapierre M, *et al.* Increased expression of the HDAC9 gene is associated with antiestrogen resistance of breast cancers. *Mol Oncol*, 2019, **13**(7): 1534-1547
- [64] Duan X, Zhang Z, Jia J, *et al.* Histone deacetylase 4 as a potential biomarker for post-stroke cognitive impairment. *Tohoku J Exp Med*, 2025, **266**(2): 153-160
- [65] Okamura M, Inoue T, Takamatsu Y, *et al.* Pharmacological inhibition of histone deacetylases ameliorates cognitive impairment after intracerebral hemorrhage with epigenetic alteration in the hippocampus. *J Stroke Cerebrovasc Dis*, 2023, **32**(9): 107275
- [66] Fang Y C, Chan L, Liou J P, *et al.* HDAC inhibitor protects chronic cerebral hypoperfusion and oxygen-glucose deprivation injuries via H3K14 and H4K5 acetylation-mediated BDNF expression. *J Cell Mol Med*, 2020, **24**(12): 6966-6977
- [67] He L, Xu W, Zhang C, *et al.* Dysregulation of vesicular glutamate transporter VGluT2 via BDNF/TrkB pathway contributes to morphine tolerance in mice. *Front Pharmacol*, 2022, **13**: 861786
- [68] Pal S, Kozono D, Yang X, *et al.* Dual HDAC and PI3K inhibition abrogates NFκB- and FOXM1-mediated DNA damage response to radiosensitize pediatric high-grade gliomas. *Cancer Res*, 2018, **78**(14): 4007-4021
- [69] Yan J, Zhang P, Tan J, *et al.* Cdk5 phosphorylation-induced SIRT2 nuclear translocation promotes the death of dopaminergic neurons in Parkinson's disease. *NPJ Parkinsons Dis*, 2022, **8**(1): 46
- [70] Hamid M A, Moustafa M T, Nashine S, *et al.* Anti-VEGF drugs influence epigenetic regulation and AMD-specific molecular markers in ARPE-19 cells. *Cells*, 2021, **10**(4): 878
- [71] Chai G S, Gong J, Wu J J, *et al.* Danggui Buxue decoction ameliorates mitochondrial biogenesis and cognitive deficits through upregulating histone H4 lysine 12 acetylation in APP/PS1 mice. *J Ethnopharmacol*, 2023, **313**: 116554
- [72] Yue P, Gao L, Wang X, *et al.* Intranasal administration of GDNF protects against neural apoptosis in a rat model of Parkinson's disease through PI3K/Akt/GSK3β pathway. *Neurochem Res*, 2017, **42**(5): 1366-1374
- [73] Soragni E, Miao W, Iudicello M, *et al.* Epigenetic therapy for friedreich ataxia. *Ann Neurol*, 2014, **76**(4): 489-508
- [74] Ma J, Krukowski K, Golonzhka O, *et al.* HDAC6 inhibition effectively reverses chemotherapy-induced peripheral neuropathy. *Brain Behav Immun*, 2017, **66**: e37
- [75] Brookes R L, Crichton S, Wolfe C D A, *et al.* Sodium valproate, a histone deacetylase inhibitor, is associated with reduced stroke risk after previous ischemic stroke or transient ischemic attack. *Stroke*, 2018, **49**(1): 54-61
- [76] Gao X, Zeb S, He Y Y, *et al.* Valproic acid inhibits glial scar formation after ischemic stroke. *Pharmacology*, 2022, **107**(5/6): 263-280
- [77] Mayerhofer E, Parodi L, Narasimhalu K, *et al.* Genetic variation supports a causal role for valproate in prevention of ischemic stroke. *medRxiv*, 2023: 2023.02.14.23285856

The Molecular Mechanisms of HDACi in Regulating Ischemic Stroke*


PAN Feng-Yuan^{1,2)}, XU Yi-Wei^{1,2,3)}, ZOU Xin-Yu^{1,2)}, CHEN Jing^{1,2)**}

⁽¹⁾College of Medicine and Health Sciences, China Three Gorges University, Yichang 443002, China;

⁽²⁾Third-grade Pharmacological Laboratory on Traditional Chinese Medicine, State Administration of Traditional Chinese Medicine, China Three Gorges University, Yichang 443002, China;

⁽³⁾Key Laboratory of Advanced Pharmaceuticals, China Three Gorges University, Yichang 443002, China)

Graphical abstract



Abstract Ischemic stroke (IS) accounts for approximately 80% of all stroke cases and is a leading cause of death and long-term disability worldwide. Its core pathological mechanism involves the interruption of cerebral blood flow, leading to neuronal cell death and ischemic tissue necrosis in the brain, which is associated with multiple molecular processes including apoptosis, inflammation, and oxidative stress. This review systematically discusses the classification of HDACs, the mechanisms of action of HDAC inhibitors, and their multiple effects in inhibiting cell apoptosis, regulating neuroinflammation, repairing the blood-brain barrier, and improving cognitive function following IS. HDACs function by removing acetyl groups from histone lysine residues, leading to chromatin condensation and gene silencing. The HDAC family is classified into four classes: class I (HDAC1, 2,

* This work was supported by a grant from Natural Science Foundation of Hubei Province (2024AFD99).

** Corresponding author.

Tel: 86-717-6392029, E-mail: chenjing@ctgu.edu.cn

Received: September 9, 2025 Accepted: November 27, 2025

3, 8), class IIa (HDAC4, 5, 7, 9), class IIb (HDAC6, 10), and class IV (HDAC11), with class III being the NAD⁺-dependent sirtuins. Histone deacetylase inhibitors (HDACi) exert significant neuroprotective effects following ischemic stroke through a multi-target, multi-pathway synergistic mechanism. The core mechanisms include inhibition of neuronal apoptosis, regulation of neuroinflammation, protection of the blood-brain barrier (BBB), and improvement of cognitive impairments (PSCI). HDACi regulate gene expression epigenetically by upregulating genes such as *p21/CIP1*, leading to cell cycle arrest, while also modulating apoptosis-related proteins by inhibiting pro-apoptotic signaling pathways, thereby reducing neuronal cell death. In terms of neuroinflammation, HDACi suppress NF- κ B and activate Nrf2 pathways, decreasing the release of pro-inflammatory cytokines and preventing the pro-inflammatory polarization of microglia and macrophages, thus modulating the inflammatory response. Regarding BBB protection, HDACi regulate the expression and restoration of tight junction proteins such as occludin and claudin-5, while inhibiting the release of destructive factors like MMP-9, alleviating vasogenic edema, and maintaining BBB integrity. Furthermore, HDACi promote the transcription of neurotrophic factors and synaptic-associated genes, enhancing neuroplasticity and repairing neuronal networks, ultimately improving cognitive functions. Therefore, HDACi demonstrate great potential as a multifaceted therapeutic strategy for ischemic stroke. HDACis represent a powerful multi-target therapeutic approach that transcends the limitations of traditional thrombolytic therapies. HDACis represent a powerful multi-target therapeutic approach that transcends the limitations of traditional thrombolytic therapies, which are hampered by a narrow time window and risks of reperfusion injury. Histone acetylation is increased by HDACis, which relaxes chromatin and reactivates protective gene transcription. Their selectivity and chemical structure are used to classify them. Trichostatin A (TSA) and sodium butyrate (SB), a short-chain fatty acid, are examples of broad-spectrum inhibitors that are effective in lowering infarct volume and reducing neuroinflammation. More selective inhibitors, including Tubastatin A (HDAC6-selective) and Entinostat (class I-selective), may have fewer adverse effects while increasing efficacy. By suppressing apoptosis by modifying the p53, Bcl-2, and JNK pathways, reducing neuroinflammation by blocking NF- κ B and NLRP3 activation, preserving the integrity of the blood-brain barrier by strengthening tight junction proteins, and promoting synaptic plasticity, neurogenesis, and the expression of neurotrophic factors like BDNF, these inhibitors provide neuroprotection through a variety of interrelated mechanisms. Despite their great potential, HDACis' clinical translation is fraught with difficulties, mostly because of non-selective inhibition-related adverse effects such as hepatotoxicity and gastrointestinal problems with valproic acid (VPA). In order to accomplish targeted delivery to the brain, future research is consequently shifting toward the development of highly selective inhibitors, refining dosing regimes, and utilizing cutting-edge drug delivery technologies like nanoparticles. In summary, the development of effective neuroprotective and neurorestorative treatments for IS may be greatly aided by a nuanced, spatiotemporally accurate understanding of HDAC activities and the judicious use of subtype-selective HDACis.

Key words histone deacetylase, histone deacetylase inhibitor, ischemic stroke, neurological protection

DOI: 10.3724/j.pibb.2025.0403

CSTR: 32369.14.pibb.20250403